

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

# Journal of Computational and Applied Mathematics

journal homepage: [www.elsevier.com/locate/cam](http://www.elsevier.com/locate/cam)

## A class of boundary value problems for first-order impulsive integro-differential equations with deviating arguments<sup>☆</sup>

Zhenhai Liu<sup>\*</sup>, Jitai Liang

College of Science, Guangxi University for Nationalities, Nanning, Guangxi 530006, PR China

### ARTICLE INFO

#### Article history:

Received 21 April 2011

Received in revised form 16 March 2012

#### Keywords:

Nonlinear impulsive integro-differential equations

Deviating arguments

Upper and lower solutions

Monotone iterative technique

### ABSTRACT

This paper is concerned with a class of boundary value problems for nonlinear mixed impulsive integro-differential equations with deviating arguments. We establish a new comparison principle and use the method of upper and lower solutions together with the monotone iterative technique. Under suitable conditions, we obtain the existence results of extremal solutions for the problems. An example is also given to illustrate our results.

© 2012 Elsevier B.V. All rights reserved.

### 1. Introduction

Impulsive differential equations have become increasingly important in recent years in some mathematical models of real processes and phenomena studied in physics, chemical technology, population dynamics, biotechnology, and economics. There has been a significant development in impulse theory. In particular, there is an increasing interest in the study of nonlinear mixed integro-differential equations with deviating arguments and multipoint boundary value problems (BVPs) [1–4] for impulsive differential equations.

In [5], the method of lower and upper solutions combined with the monotone iterative technique and the numerical-analytic method were applied to study the problem

$$\begin{cases} x'(t) = f\left(t, x(t), \int_0^T k(s)x(s)ds\right) & t \in J = [0, T] \\ x(0) = \lambda x(T) + \int_0^T D(s)x(s)ds + d & d \in R, \end{cases}$$

where  $f \in C[J \times R^2, R]$ ,  $f$  is non-decreasing with respect to the third variable,  $k, D \in C[J, R_+]$ , and  $\lambda \geq 0$ .

Chen and Shen [6] studied

$$\begin{cases} u'(t) = f(t, u(t), u(\theta(t))) & t \neq t_k, t \in J = [0, T] \\ \Delta u(t_k) = I_k(u(t_k)) & k = 1, 2, \dots, m \\ u(0) + \mu \int_0^T u(s)ds = u(T), \end{cases}$$

where  $f \in C[J \times R^2, R]$ , and  $\mu = 1$  or  $-1$ , by the method of upper and lower solutions and the monotone iterative technique.

<sup>☆</sup> Project supported by NNSF of China Grant No. 10971019 and the Innovation Project of Guangxi University for Nationalities (gxun-chx2011080).

<sup>\*</sup> Corresponding author. Tel.: +86 771 3265663; fax: +86 771 3261718.

E-mail addresses: [zhhlui100@yahoo.com.cn](mailto:zhhlui100@yahoo.com.cn), [zhhlui@hotmail.com](mailto:zhhlui@hotmail.com), [ldolmy@163.com](mailto:ldolmy@163.com) (Z. Liu).

Motivated by the above, we are concerned with the following BVPs of nonlinear mixed impulsive integro-differential equations with deviating arguments:

$$\begin{cases} u'(t) = f(t, u(t), u(\alpha(t)), Tu, Su) & t \neq t_k, t \in J = [0, T] \\ \Delta u(t_k) = I_k(u(t_k)) & k = 1, 2, \dots, m \\ u(0) = \lambda_1 u(T) + \lambda_2 u(\eta) + \lambda_3 \int_0^T w(s, u(s)) ds + k, \end{cases} \quad (1.1)$$

where  $0 = t_0 < t_1 < t_2 < \dots < t_k < \dots < t_m < t_{m+1} = T$ ,  $I_k \in C(R, R)$ ,  $f$  is continuous everywhere except at  $\{t_k\} \times R^4$ ;  $f(t_k^+, \cdot, \cdot, \cdot, \cdot)$  and  $f(t_k^-, \cdot, \cdot, \cdot, \cdot)$  exist with  $f(t_k^-, \cdot, \cdot, \cdot, \cdot) = f(t_k, \cdot, \cdot, \cdot, \cdot)$ ;

$$(Tu)(t) = \int_0^{\beta(t)} k(t, s)u(\gamma(s))ds, \quad (Su)(t) = \int_0^T h(t, s)u(\delta(s))ds,$$

and  $\Delta u(t_k) = u(t_k^+) - u(t_k^-)$ ,  $w \in C(J \times R, R)$ ,  $0 \leq \lambda_1 \leq 1$ ,  $0 \leq \lambda_2$ ,  $0 \leq \lambda_3$ ,  $k \in R$ , and  $0 \leq \eta \leq T$ . The assumptions concerning  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ ,  $k$ , and  $h$  will be given latter. The boundary conditions in Eq. (1.1) involve several special cases such as periodic boundary conditions, anti-periodic boundary conditions, integral boundary, and initial problems.

*Special cases*

- (i) If  $\lambda_1 = 1$ ,  $\lambda_2 = \lambda_3 = k = 0$ , then Eq. (1.1) reduces to the periodic boundary value problem (cf. [7–11]).
- (ii) If  $\lambda_2 = 1 + \lambda_1$ ,  $\eta = 0$ , and  $\lambda_3 = k = 0$ , then Eq. (1.1) reduces to the anti-periodic boundary value problem (cf. [12–15]).
- (iii) If  $\lambda_3 \neq 0$  and  $\lambda_2 = k = 0$ , then Eq. (1.1) reduces to the integral boundary value problems which have been studied in [16–19].
- (iv) If  $\lambda_1 = \lambda_2 = \lambda_3 = 0$ , then Eq. (1.1) reduces to initial problems (cf. [20,21]).

For example, if  $\lambda_2 = 1 + \lambda_1$ ,  $\eta = 0$ , and  $\lambda_3 = k = 0$ , Eq. (1.1) reduces to an anti-periodic boundary value problem, which was considered by Wang and Zhang in [15]. There, the existence results of quasi-extremal solutions for the anti-periodic boundary value problems was obtained by the method of upper and lower solutions with the monotone iterative technique. Therefore, we extend some previous results in many respects.

The article is organized as follow. In Section 2, we establish a new comparison principle. In Section 3, by using of the monotone iterative technique and the method of upper and lower solutions, we obtain the existence results of extremal solutions for (1.1). In Section 4, we give an example that illustrates our results.

## 2. Preliminaries and lemmas

Let  $PC(J) = \{x : J \rightarrow R; x(t) \text{ is continuous everywhere except for some } t_k \text{ at which } x(t_k^+) \text{ and } x(t_k^-) \text{ exist and } x(t_k) = x(t_k^-), k = 1, 2, \dots, m\}$ ;  $PC^1(J) = \{x \in PC(J) : x'(t) \text{ is continuous everywhere except for some } t_k \text{ at which } x'(t_k^+) \text{ and } x'(t_k^-) \text{ exist and } x'(t_k) = x'(t_k^-), k = 1, 2, \dots, m\}$ . Let  $J^- = J \setminus \{t_k, k = 1, 2, \dots, m\}$ ;  $PC(J)$  and  $PC^1(J)$  are Banach spaces with the norms  $\|x\|_{PC} = \sup\{|x(t)| : t \in J\}$  and  $\|x\|_{PC^1} = \max\{\|x\|_{PC}, \|x'\|_{PC}\}$ .  $x \in PC^1(J)$  is called a solution of BVPs (1.1) if it satisfies Eq. (1.1).

In what follows, we need the following hypotheses.

- (H<sub>1</sub>)  $\alpha, \beta, \gamma, \delta \in C(J, J)$ ,  $N, K, H \in C(J, R_+)$ ,  $k \in C(\Omega, R_+)$ ,  $h \in C(J^2, R_+)$ ,  $R_+ = [0, +\infty)$ ,  $\Omega = \{(t, s) \in J^2 \mid 0 \leq s \leq \beta(t)\}$ ,  $M \in C(J, R)$ ,  $\int_0^T M(\tau) d\tau \geq 0$ , and  $0 \leq L_k \leq 1$ ,  $0 < r \leq 1$ .

For convenience, we set

$$\begin{cases} N^*(t) = N(t)e^{\int_0^t M(s)ds}e^{-\int_0^{\alpha(t)} M(s)ds}, & K^*(t) = K(t)e^{\int_0^t M(s)ds}, \\ H^*(t) = H(t)e^{\int_0^t M(s)ds}, & k^*(t, s) = k(t, s)e^{-\int_0^{\gamma(s)} M(\tau)d\tau}, \\ h^*(t, s) = h(t, s)e^{-\int_0^{\delta(s)} M(\tau)d\tau}, & r^* = re^{-\int_0^T M(s)ds}, \end{cases} \quad (2.1)$$

$$(H_2) \quad \theta^*(t) = N^*(t) + K^*(t) \int_0^{\beta(t)} k^*(t, s)ds + H^*(t) \int_0^T h^*(t, s)ds \neq 0 \text{ for } t \in J, \mu^* = \int_0^T \theta^*(t)dt.$$

$$\left[ \mu^* + \sum_{k=1}^m L_k \right] \leq r^*.$$

**Lemma 2.1.** Assume that (H<sub>1</sub>) and (H<sub>2</sub>) hold, and that  $q \in PC^1(J)$  such that

$$\begin{cases} q'(t) \leq -M(t)q(t) - (\mathcal{H}q)(t) & t \neq t_k, t \in J = [0, T] \\ \Delta q(t_k) \leq -L_k(q(t_k)) & k = 1, 2, \dots, m \\ q(0) \leq rq(T), \end{cases} \quad (2.2)$$

where the operator  $\mathcal{H}$  is defined as

$$(\mathcal{H}q)(t) = N(t)q(\alpha(t)) + K(t) \int_0^{\beta(t)} k(t, s)q(\gamma(s))ds + H(t) \int_0^T h(t, s)q(\delta(s))ds.$$

Then  $q(t) \leq 0$  for  $t \in J$ .

**Proof.** Let  $p(t) = q(t)e^{\int_0^t M(s)ds}$ . Obviously  $p(t)$  and  $q(t)$  have the same sign on  $J$ . In view of (2.2), we have

$$\begin{cases} p'(t) \leq -(\mathcal{H}^*p)(t) & t \neq t_k, \quad t \in J = [0, T] \\ \Delta p(t_k) \leq -L_k(p(t_k)) & k = 1, 2, \dots, m \\ p(0) \leq r^*p(T), \end{cases} \quad (2.3)$$

where  $(\mathcal{H}^*p)(t) = N^*(t)p(\alpha(t)) + K^*(t) \int_0^{\beta(t)} k^*(t, s)p(\gamma(s))ds + H^*(t) \int_0^T h^*(t, s)p(\delta(s))ds$ .

Next, we will show that  $p(t) \leq 0$ .

Suppose, to the contrary, that  $p(t) > 0$  for some  $t \in J$ .

(i) If  $p(t) \geq 0$ ,  $p(t) \not\equiv 0$  for  $t \in J$ , we get  $p'(t) \leq 0$ , in view of the first inequality of (2.3). By the second inequality in (2.3), we obtain that  $p(t)$  is non-increasing in  $J$ . Then  $0 \leq p(T) \leq p(t) \leq p(0)$ . On the other hand, by the third inequality in (2.3), if  $r^* = 1$ , then  $p(T) \leq p(t) \leq p(0) \leq p(T)$ , so we get that  $p(t) \equiv C > 0$ . Hence  $p'(t) \equiv 0$ . By the first inequality in (2.3) again, we have

$$0 \leq -C\theta^*(t) \quad \forall t \in J.$$

By  $(H_1)$ , we get that  $C \leq 0$ , which is a contradiction.

If  $0 < r^* < 1$ , then  $p(T) \leq p(0) \leq r^*p(T)$ , so  $p(T)(1 - r^*) \leq 0$ . We have  $0 \leq p(T) \leq 0$ . Since  $p$  is non-increasing in  $J$ , we infer that  $p(t) \equiv 0$ . This is a contradiction.

(ii) If  $p(t^*) = \sup_{t \in J} p(t) > 0$ ,  $p(t_*) = \inf_{t \in J} p(t) = -\lambda < 0$ , then  $\lambda > 0$ .

Case 1. If  $t_* < t^*$ , integrating from  $t_*$  to  $t^*$ , we get, from (2.3), that

$$\begin{aligned} 0 < p(t^*) &= p(t_*) + \int_{t_*}^{t^*} p'(s)ds + \sum_{t_* \leq t_k < t^*} \Delta p(t_k) \\ &\leq -\lambda + \int_{t_*}^{t^*} -(\mathcal{H}^*p)(s)ds - \sum_{t_* \leq t_k < t^*} L_k p(t_k) \\ &\leq -\lambda + \mu^* \lambda + \lambda \sum_{k=1}^m L_k. \end{aligned}$$

Hence

$$1 < \mu^* + \sum_{k=1}^m L_k,$$

which is in contradiction to  $(H_2)$ .

Case 2. If  $t^* < t_*$ , we have

$$\begin{aligned} 0 < p(t^*) &= p(0) + \int_0^{t^*} p'(s)ds + \sum_{0 < t_k < t^*} \Delta p(t_k) \\ &\leq p(0) + \int_0^{t^*} -(\mathcal{H}^*p)(s)ds + \lambda \sum_{0 < t_k < t^*} L_k \\ &\leq p(0) + \lambda \int_0^{t^*} \theta^*(s)ds + \lambda \sum_{0 < t_k < t^*} L_k, \\ p(T) &= p(t_*) + \int_{t_*}^T p'(s)ds + \sum_{t_* \leq t_k < T} \Delta p(t_k) \\ &\leq -\lambda + \int_{t_*}^T -(\mathcal{H}^*p)(s)ds + \lambda \sum_{t_* \leq t_k < T} L_k \\ &\leq -\lambda + \lambda \int_{t_*}^T \theta^*(s)ds + \lambda \sum_{t_* \leq t_k < T} L_k. \end{aligned}$$

By the two inequalities above, we obtain

$$\begin{aligned}
 -\lambda + \frac{1}{r^*} \lambda \int_{t_*}^T \theta^*(s) ds + \frac{1}{r^*} \lambda \sum_{t_* \leq t_k < T} L_k &\geq -\lambda + \lambda \int_{t_*}^T \theta^*(s) ds + \lambda \sum_{t_* \leq t_k < T} L_k \\
 &\geq p(T) \geq \frac{1}{r^*} p(0) \\
 &> -\frac{1}{r^*} \lambda \int_0^{t_*} \theta^*(s) ds - \frac{1}{r^*} \lambda \sum_{0 < t_k < t_*} L_k \\
 &\geq -\frac{1}{r^*} \lambda \int_0^{t_*} \theta^*(s) ds - \frac{1}{r^*} \lambda \sum_{0 < t_k < t_*} L_k.
 \end{aligned}$$

Therefore, we get that  $(\mu^* + \sum_{k=1}^m L_k) > r^*$ , which is in contradiction to  $(H_2)$ . Hence  $p(t) \leq 0$ ,  $q(t) \leq 0$ . We complete the proof.  $\square$

Let  $C_k, d \in R, \sigma \in PC(J)$ . In order to deal with the following linear problem,

$$\begin{cases} u'(t) = -M(t)u(t) - (\mathcal{H}u)(t) + \sigma(t) & t \neq t_k, t \in J = [0, T] \\ \Delta u(t_k) = -L_k(u(t_k)) + C_k & k = 1, 2, \dots, m \\ u(0) = ru(T) + d, \end{cases} \quad (2.4)$$

we also need the following hypothesis:

$$(H_3) \quad \varpi \equiv e^{\int_0^T |M(\tau)| d\tau} \left( 1 + \frac{r}{e^{\int_0^T M(\tau) d\tau} - r} \right) \left( \mu + \sum_{k=1}^m L_k \right) < 1, \quad (2.5)$$

where  $\mu = \int_0^T [N(t) + K(t) \int_0^{\beta(t)} k(t, s) ds + H(t) \int_0^T h(t, s) ds] dt$ .

**Lemma 2.2.** Suppose that  $(H_3)$  holds and that  $0 \leq L_k \leq 1, M \in C(J, R), \int_0^T M(s) ds \geq 0, 0 < r \leq 1$ , and  $\int_0^T M(s) ds > 0$  if  $r = 1$ . Then the following integral equation,

$$\begin{aligned}
 u(t) &= \frac{de^{\int_0^T M(\tau) d\tau}}{e^{\int_0^T M(\tau) d\tau} - r} + \int_0^T G(t, s)(\sigma(s) - (\mathcal{H}u)(s)) ds + \frac{re^{-\int_0^t M(\tau) d\tau}}{e^{\int_0^T M(\tau) d\tau} - r} \sum_{k=1}^m e^{\int_0^{t_k} M(\tau) d\tau} (-L_k(u(t_k)) + C_k) \\
 &+ \sum_{0 < t_k < t} e^{-\int_0^t M(\tau) d\tau} e^{\int_0^{t_k} M(\tau) d\tau} (-L_k(u(t_k)) + C_k), \quad (2.6)
 \end{aligned}$$

where

$$G(t, s) = \begin{cases} \frac{e^{\int_t^T M(\tau) d\tau} e^{\int_0^s M(\tau) d\tau}}{e^{\int_0^T M(\tau) d\tau} - r}, & 0 \leq s \leq t \leq T, \\ \frac{re^{\int_t^s M(\tau) d\tau}}{e^{\int_0^T M(\tau) d\tau} - r}, & 0 \leq t \leq s \leq T, \end{cases}$$

has a unique solution  $u$  in  $PC(J)$ .

**Proof.** Define operator  $F$  by

$$\begin{aligned}
 (Fu)(t) &= \frac{de^{\int_0^T M(\tau) d\tau}}{e^{\int_0^T M(\tau) d\tau} - r} + \int_0^T G(t, s)(\sigma(s) - (\mathcal{H}u)(s)) ds \\
 &+ \frac{re^{-\int_0^t M(\tau) d\tau}}{e^{\int_0^T M(\tau) d\tau} - r} \sum_{k=1}^m e^{\int_0^{t_k} M(\tau) d\tau} (-L_k(u(t_k)) + C_k) + \sum_{0 < t_k < t} e^{-\int_0^t M(\tau) d\tau} e^{\int_0^{t_k} M(\tau) d\tau} (-L_k(u(t_k)) + C_k).
 \end{aligned}$$

If  $0 \leq s \leq t \leq T$ ,

$$\begin{aligned}
 \frac{e^{\int_t^T M(\tau) d\tau} e^{\int_0^s M(\tau) d\tau}}{e^{\int_0^T M(\tau) d\tau} - r} &\leq \frac{e^{\int_0^T M(\tau) d\tau}}{e^{\int_0^T M(\tau) d\tau} - r} \\
 &= \frac{e^{\int_0^T M(\tau) d\tau} - r + r}{e^{\int_0^T M(\tau) d\tau} - r} \\
 &= 1 + \frac{r}{e^{\int_0^T M(\tau) d\tau} - r},
 \end{aligned}$$

and if  $0 \leq t \leq s \leq T$ ,

$$\begin{aligned} \frac{re^{\int_t^s M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} &\leq \frac{re^{\int_0^T M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} \\ &\leq \frac{e^{\int_0^T M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} \\ &= 1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r}, \end{aligned}$$

so it is easy to see that

$$\max\{G(t, s), (t, s) \in J^2\} = 1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r}.$$

For any  $x, y \in PC(J)$ , we get

$$\begin{aligned} \|(Fx)(t) - (Fy)(t)\|_{PC} &\leq \left(1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r}\right) \int_0^T |(-\mathcal{H}x(s) + \mathcal{H}y(s))ds| \\ &\quad + \max \left| \frac{re^{-\int_0^t M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} \sum_{k=1}^m e^{\int_0^{t_k} M(\tau)d\tau} (-L_k(x(t_k)) + L_k(y(t_k))) \right. \\ &\quad \left. + \sum_{0 < t_k < t} e^{-\int_0^t M(\tau)d\tau} e^{\int_0^{t_k} M(\tau)d\tau} (-L_k(x(t_k)) + L_k(y(t_k))) \right| \\ &\leq \left(1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r}\right) \int_0^T |(-\mathcal{H}x(s) + \mathcal{H}y(s))ds| \\ &\quad + \max \left\{ \left(1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r}\right) \sum_{0 < t_k < t} e^{-\int_0^{t_k} M(\tau)d\tau} |(-L_k(x(t_k)) + L_k(y(t_k)))| \right. \\ &\quad \left. + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r} \sum_{t \leq t_k < T} e^{\int_t^{t_k} M(\tau)d\tau} |(-L_k(x(t_k)) + L_k(y(t_k)))| \right\} \\ &\leq \left(1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r}\right) \int_0^T |(-\mathcal{H}x(s) + \mathcal{H}y(s))ds| \\ &\quad + \left(1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r}\right) \max \left\{ \sum_{0 < t_k < t} e^{-\int_0^{t_k} M(\tau)d\tau} |(-L_k(x(t_k)) + L_k(y(t_k)))| \right. \\ &\quad \left. + \sum_{t \leq t_k < T} e^{\int_t^{t_k} M(\tau)d\tau} |(-L_k(x(t_k)) + L_k(y(t_k)))| \right\} \\ &\leq \left(1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r}\right) \int_0^T |(-\mathcal{H}x(s) + \mathcal{H}y(s))ds| \\ &\quad + \left(1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r}\right) \max \left\{ \sum_{0 < t_k < t} e^{\int_0^T |M(\tau)|d\tau} |(-L_k(x(t_k)) + L_k(y(t_k)))| \right. \\ &\quad \left. + \sum_{t \leq t_k < T} e^{\int_0^T |M(\tau)|d\tau} |(-L_k(x(t_k)) + L_k(y(t_k)))| \right\} \\ &\leq e^{\int_0^T |M(\tau)|d\tau} \left(1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r}\right) \left(\mu + \sum_{k=1}^m L_k\right) \|x - y\|_{PC} \\ &= \varpi \|x - y\|_{PC}, \end{aligned}$$

which implies from the Banach fixed point theorem that  $F$  has a unique fixed point  $u$  in  $PC(J)$ . The proof is complete.  $\square$

**Lemma 2.3.** Assume that  $(H_1)$ – $(H_3)$  hold and that  $\int_0^T M(s)ds > 0$  if  $r = 1$ . Then the linear problem (2.4) has a unique solution  $u \in PC^1(J, E)$ , and it is represented by the integral equation (2.6).

**Proof.** By Lemma 2.2, the integral equation (2.6) has a unique solution  $u \in PC(J)$ . Differentiating (2.6), we obtain

$$\begin{aligned} u'(t) &= \frac{d}{dt} \left[ \frac{de^{\int_0^t M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} + \int_0^T G(t, s)(\sigma(s) - (\mathcal{H}u)(s))ds + \frac{re^{-\int_0^t M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} \right. \\ &\quad \times \sum_{k=1}^m e^{\int_0^{t_k} M(\tau)d\tau} (-L_k(u(t_k)) + C_k) + \sum_{0 < t_k < t} e^{-\int_0^t M(\tau)d\tau} e^{\int_0^{t_k} M(\tau)d\tau} (-L_k(u(t_k)) + C_k) \left. \right] \\ &= -M(t) \left[ \frac{de^{\int_0^t M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} + \int_0^T G(t, s)(\sigma(s) - (\mathcal{H}u)(s))ds + \frac{re^{-\int_0^t M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} \right. \\ &\quad \times \sum_{k=1}^m e^{\int_0^{t_k} M(\tau)d\tau} (-L_k(u(t_k)) + C_k) + \sum_{0 < t_k < t} e^{-\int_0^t M(\tau)d\tau} e^{\int_0^{t_k} M(\tau)d\tau} (-L_k(u(t_k)) + C_k) \left. \right] \\ &\quad + \left( \frac{-r}{e^{\int_0^T M(\tau)d\tau} - r} + \frac{e^{\int_0^T M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} \right) (\sigma(t) - (\mathcal{H}u)(t)) \\ &= -M(t)u(t) - (\mathcal{H}u)(t) + \sigma(t) \quad t \in J^-, \\ \Delta u(t_k) &= u(t_k^+) - u(t_k^-) \\ &= \sum_{0 < t_j \leq t_k} \Delta u(t_j) - \sum_{0 < t_j < t_k} \Delta u(t_j) \\ &= \sum_{j=1}^k (-L_j(u(t_j)) + C_j) - \sum_{j=1}^{k-1} (-L_j(u(t_j)) + C_j) \\ &= -L_k(u(t_k)) + C_k. \end{aligned}$$

Also,

$$\begin{aligned} u(0) &= \frac{r}{e^{\int_0^T M(\tau)d\tau} - r} \sum_{k=1}^m e^{\int_0^{t_k} M(\tau)d\tau} (-L_k(u(t_k)) + C_k) \\ &\quad + \int_0^T \frac{re^{\int_0^s M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} (\sigma(s) - (\mathcal{H}u)(s))ds + \frac{de^{\int_0^T M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r}, \\ u(T) &= \frac{1}{e^{\int_0^T M(\tau)d\tau} - r} \sum_{k=1}^m e^{\int_0^{t_k} M(\tau)d\tau} (-L_k(u(t_k)) + C_k) \\ &\quad + \int_0^T \frac{e^{\int_0^s M(\tau)d\tau}}{e^{\int_0^T M(\tau)d\tau} - r} (\sigma(s) - (\mathcal{H}u)(s))ds + \frac{d}{e^{\int_0^T M(\tau)d\tau} - r}. \end{aligned}$$

It is easy to check that  $u(0) = ru(T) + d$ .

Hence, we infer that  $u$  defined by (2.6) is a solution of (2.4).

Next, we show that problem (2.4) is uniquely solvable. Let  $u_1, u_2$  be two solutions of (2.4) and set  $p = u_1 - u_2$ . We have

$$\begin{aligned} p' &= u_1' - u_2' \\ &= -M(t)u_1(t) - (\mathcal{H}u_1)(t) + \sigma(t) - (-M(t)u_2(t) - (\mathcal{H}u_2)(t) + \sigma(t)) \\ &= -M(t)p(t) - (\mathcal{H}p)(t) \quad t \in J^-, \\ \Delta p(t_k) &= \Delta u_1 - \Delta u_2 \\ &= -L_k u_1(t_k) + C_k - (-L_k u_2(t_k) + C_k) \\ &= -L_k p(t_k), \\ p(0) &= u_1(0) - u_2(0) \\ &= ru_1(T) + d - (ru_2(T) + d) \\ &= rp(T). \end{aligned}$$

In view of Lemma 2.1, we get that  $p \leq 0$ , which implies that  $u_1 \leq u_2$ . Similarly, we can get that  $u_1 \geq u_2$ . Hence  $u_1 = u_2$ . The proof is complete.  $\square$

### 3. Main results

If  $u_0, v_0 \in PC^1(J)$  and  $u_0(t) \leq v_0(t)$ ,  $\forall t \in J$ , we define the interval

$$[u_0, v_0] = \{x \in PC^1(J) : u_0(t) \leq x(t) \leq v_0(t), t \in J\}.$$

In what follows, we also need the following assumptions.

(H<sub>4</sub>) There exist  $u_0, v_0 \in PC^1(J)$  with  $u_0(t) \leq v_0(t)$ ,  $\forall t \in J$  such that

$$\begin{cases} u'_0(t) \leq f(t, u_0(t), u_0(\alpha(t)), Tu_0, Su_0) & t \neq t_k, t \in J = [0, T] \\ \Delta u_0(t_k) \leq I_k(u_0(t_k)) & k = 1, 2, \dots, m \\ u_0(0) \leq \lambda_1 u_0(T) + \lambda_2 u_0(\eta) + \lambda_3 \int_0^T w(s, u_0(s))ds + k, \\ v'_0(t) \geq f(t, v_0(t), v_0(\alpha(t)), Tv_0, Sv_0) & t \neq t_k, t \in J = [0, T] \\ \Delta v_0(t_k) \geq I_k(v_0(t_k)) & k = 1, 2, \dots, m \\ v_0(0) \geq \lambda_1 v_0(T) + \lambda_2 v_0(\eta) + \lambda_3 \int_0^T w(s, v_0(s))ds + k. \end{cases} \quad (3.1)$$

(H<sub>5</sub>)

$$f(t, \bar{u}, \bar{u}(\alpha(t)), T\bar{u}, S\bar{u}) - f(t, u, u(\alpha(t)), Tu, Su) \geq -M(t)(\bar{u} - u) - N(t)(\bar{u} - u)(\alpha(t)) \\ - K(t)T(\bar{u} - u) - H(t)S(\bar{u} - u), \quad (3.2)$$

$$I_k(\bar{u}) - I_k(u) \geq -L_k(\bar{u} - u), \quad (3.3)$$

for all  $u_0(t) \leq u(t) \leq \bar{u}(t) \leq v_0(t)$  in  $J$ .

(H<sub>6</sub>) Assume that  $a(t)$  is a non-negative integrable function, such that

$$w(t, \bar{u}) - w(t, u) \geq a(t)(\bar{u} - u), \quad (3.4)$$

for all  $u_0(t) \leq u(t) \leq \bar{u}(t) \leq v_0(t)$  in  $J$ .

**Theorem 3.1.** Assume that (H<sub>1</sub>)–(H<sub>6</sub>) hold and that  $\int_0^T M(s)ds > 0$  if  $r = 1$ . Then there exist two monotone sequences  $\{u_n(t)\}, \{v_n(t)\} \subset PC^1(J)$  with

$$u_0 \leq u_1 \leq \dots \leq u_n \leq \dots \leq v_n \leq \dots \leq v_1 \leq v_0, \quad (3.5)$$

such that  $\lim_{n \rightarrow \infty} u_n = u^*(t)$ ,  $\lim_{n \rightarrow \infty} v_n = v^*(t)$  uniformly on  $J$ . Moreover,  $u^*(t)$  and  $v^*(t)$  are the minimal solution and the maximal solution of (1.1) in  $[u_0, v_0]$ , respectively.

**Proof.** For  $\xi \in [u_0, v_0]$ , we consider (2.4) with

$$\sigma(t) = M(t)\xi(t) + (\mathcal{H}\xi)(t) + f(t, \xi(t), \xi(\alpha(t)), T\xi, S\xi), \quad C_k = I_k(\xi(t_k)) + L_k\xi(t_k), \\ d = \lambda_2\xi(\eta) + \lambda_3 \int_0^T w(s, \xi(s))ds + k, \quad r = \lambda_1.$$

By Lemma 2.3, the BVP (2.4) has a unique solution  $u \in PC(J)$ .

We define an operator  $A : [u_0, v_0] \rightarrow PC(J)$  by  $u = A\xi$ . We claim that

(a)  $u_0 \leq Au_0, Av_0 \leq v_0$ ,

(b)  $A$  is nondecreasing on  $[u_0, v_0]$ .

We first prove (a). Set  $u_1 = Au_0$ ,  $p(t) = u_0(t) - u_1(t)$ . Then we have

$$\begin{aligned} p' &= u'_0 - u'_1 \\ &\leq f(t, u_0(t), u_0(\alpha(t)), Tu_0, Su_0) - [f(t, u_0(t), u_0(\alpha(t)), Tu_0, Su_0) \\ &\quad + M(t)u_0(t) + (\mathcal{H}u_0)(t) - M(t)u_1(t) - (\mathcal{H}u_1)(t)] \\ &= -M(t)p(t) - (\mathcal{H}p)(t) \quad t \in J^-, \\ \Delta p(t_k) &= \Delta u_0(t_k) - \Delta u_1(t_k) \\ &\leq I_k(u_0(t_k)) - [I_k(u_0(t_k)) - L_k(u_1 - u_0)] \\ &= -L_k p(t_k), \\ p(0) &= u_0(0) - u_1(0) \\ &\leq \lambda_1 u_0(T) + \lambda_2 u_0(\eta) + \lambda_3 \int_0^T w(s, u_0(s))ds + k - (\lambda_1 u_1(T) + \lambda_2 u_0(\eta) + \lambda_3 \int_0^T w(s, u_0(s))ds + k) \\ &= \lambda_1 p(T). \end{aligned}$$

By Lemma 2.1, we have  $p \leq 0$ . That is,  $u_0 \leq Au_0$ . Similarly, we can prove that  $Av_0 \leq v_0$ .

To prove (b), set  $\gamma_1, \gamma_2 \in [u_0, v_0]$  and  $\gamma_1 \leq \gamma_2$ ,  $\gamma_1^* = A\gamma_1$ ,  $\gamma_2^* = A\gamma_2$ ,  $p = \gamma_1^* - \gamma_2^*$ . Then

$$\begin{aligned} p'(t) &= \gamma_1'^* - \gamma_2'^* \\ &= f(t, \gamma_1(t), \gamma_1(\alpha(t)), T\gamma_1, S\gamma_1) + M(t)\gamma_1(t) + (\mathcal{H}\gamma_1)(t) - M(t)\gamma_1^*(t) - (\mathcal{H}\gamma_1^*)(t) \\ &\quad - [f(t, \gamma_2(t), \gamma_2(\alpha(t)), T\gamma_2, S\gamma_2) + M(t)\gamma_2(t) + (\mathcal{H}\gamma_2)(t) - M(t)\gamma_2^*(t) - (\mathcal{H}\gamma_2^*)(t)] \\ &\leq -M(t)p(t) - (\mathcal{H}p)(t) \quad t \in J^-, \\ \Delta p(t_k) &= \Delta\gamma_1^*(t_k) - \Delta\gamma_2^*(t_k) \\ &= I_k(\gamma_1(t_k)) - L_k(\gamma_1^*(t_k) - \gamma_1(t_k)) - (I_k(\gamma_2(t_k)) - L_k(\gamma_2^*(t_k) - \gamma_2(t_k))) \\ &= I_k(\gamma_1(t_k)) - I_k(\gamma_2(t_k)) + L_k(\gamma_1 - \gamma_2) - L_k(\gamma_1^* - \gamma_2^*) \\ &\leq -L_k p(t_k), \\ p(0) &= \gamma_1^*(0) - \gamma_2^*(0) \\ &\leq \lambda_1 \gamma_1^*(T) + \lambda_2 \gamma_1(\eta) + \lambda_3 \int_0^T w(s, \gamma_1(s))ds + k - (\lambda_1 \gamma_2^*(T) + \lambda_2 \gamma_2(\eta) + \lambda_3 \int_0^T w(s, \gamma_2(s))ds + k) \\ &\leq \lambda_1 p(T) + \lambda_2 (\gamma_1(\eta) - \gamma_2(\eta)) + \lambda_3 \int_0^T a(s)(\gamma_1(s) - \gamma_2(s))ds \\ &\leq \lambda_1 p(T). \end{aligned}$$

In view of Lemma 2.1, we get that  $A\gamma_1 \leq A\gamma_2$ . Hence (b) holds.

Now, we define two sequences  $\{u_n\}, \{v_n\}$  in  $PC^1(J)$ :

$$u_{n+1} = Au_n, \quad v_{n+1} = Av_n \quad (n = 0, 1, 2, \dots).$$

By (a) and (b), we have that (3.5) holds. And, for any  $n = 1, 2, \dots, u_n$  and  $v_n \in PC^1(J)$  satisfy the following integro-differential equations:

$$\begin{cases} u_n'(t) = f(t, u_{n-1}(t), u_{n-1}(\alpha(t)), Tu_{n-1}, Su_{n-1}) - M(t)(u_n(t) - u_{n-1}(t)) \\ \quad - (\mathcal{H}(u_n - u_{n-1}))(t) \quad t \neq t_k, \quad t \in J = [0, T] \\ \Delta u_n(t_k) = -L_k u_n(t_k) + I_k(u_{n-1}(t_k)) + L_k u_{n-1}(t_k) \quad k = 1, 2, \dots, m \\ u_n(0) = \lambda_1 u_n(T) + \lambda_2 u_{n-1}(\eta) + \lambda_3 \int_0^T w(s, u_{n-1}(s))ds + k, \\ v_n'(t) = f(t, v_{n-1}(t), v_{n-1}(\alpha(t)), Tv_{n-1}, Sv_{n-1}) - M(t)(v_n(t) - v_{n-1}(t)) \\ \quad - (\mathcal{H}(v_n - v_{n-1}))(t) \quad t \neq t_k, \quad t \in J = [0, T] \\ \Delta v_n(t_k) = -L_k v_n(t_k) + I_k(v_{n-1}(t_k)) + L_k v_{n-1}(t_k) \quad k = 1, 2, \dots, m \\ v_n(0) = \lambda_1 v_n(T) + \lambda_2 v_{n-1}(\eta) + \lambda_3 \int_0^T w(s, v_{n-1}(s))ds + k. \end{cases}$$

Therefore, we have that  $\{u_n\}, \{v_n\}$  are monotonically and uniformly convergent to  $u^*(t)$  and  $v^*(t)$  on  $J$ , respectively. It is not difficult to prove that  $u^*(t), v^*(t)$  are solutions of Eq. (1.1).

Finally, we assert that, if  $u \in [u_0, v_0]$  is any solution of (1.1), then  $u^*(t) \leq u(t) \leq v^*(t)$  on  $J$ . We will prove that, if  $u_n \leq u \leq v_n$ , for  $n = 0, 1, 2, \dots$ , then  $u_{n+1}(t) \leq u(t) \leq v_{n+1}(t)$ .

Set  $p(t) = u_{n+1}(t) - u(t)$ . Then

$$\begin{aligned} p'(t) &= u_{n+1}' - u'(t) \\ &= f(t, u_n(t), Tu_n, Su_n) + M(t)u_n(t) + (\mathcal{H}u_n)(t) \\ &\quad - M(t)u_{n+1}(t) - (\mathcal{H}u_{n+1})(t) - f(t, u(t), u(\alpha(t)), Tu, Su) \\ &\leq -M(t)(u_{n+1}(t) - u(t)) - (\mathcal{H}(u_{n+1} - u))(t) \\ &\leq -M(t)p(t) - (\mathcal{H}p)(t) \quad t \in J^-, \\ \Delta p(t_k) &= \Delta u_{n+1}(t_k) - \Delta u(t_k) \\ &= I_k(u_n(t_k)) - L_k(u_{n+1}(t_k) - u_n(t_k)) - I_k u(t_k) \\ &\leq -L_k(u_n(t_k) - u(t_k)) - L_k(u_{n+1}(t_k) - u_n(t_k)) \\ &= -L_k(u_{n+1}(t_k) - u(t_k)) \\ &= -L_k p(t_k), \\ p(0) &= u_{n+1}(0) - u(0) \\ &\leq \lambda_1 u_{n+1}(T) + \lambda_2 u_n(\eta) + \lambda_3 \int_0^T w(s, u_n(s))ds + k - (\lambda_1 u(T) + \lambda_2 u(\eta) + \lambda_3 \int_0^T w(s, u(s))ds + k) \\ &\leq \lambda_1 p(T) + \lambda_2 (u_n(\eta) - u(\eta)) + \lambda_3 \int_0^T a(s)(u_n(s) - u(s))ds \\ &\leq \lambda_1 p(T). \end{aligned}$$



By Lemma 2.1, we have that  $p(t) \leq 0$  for all  $t \in J$ . That is,  $u_{n+1}(t) \leq u(t)$ . Similarly, we can prove that  $u(t) \leq v_{n+1}(t)$  for all  $t \in J$ . Thus  $u_{n+1}(t) \leq u(t) \leq v_{n+1}(t)$  for all  $t \in J$ , which implies that  $u^*(t) \leq u(t) \leq v^*(t)$ . The proof is complete.  $\square$

#### 4. Example

Consider the following problems:

$$\begin{cases} u'(t) = \frac{t^4 u(t)}{100} - \frac{t}{600} \sin\left(u\left(\frac{t}{2}\right)\right) - \frac{t}{100} \int_0^t su(s)ds - \frac{t^3}{1000} \int_0^1 u(s)ds & t \neq \frac{1}{2}, t \in J = [0, 1] \\ \Delta u\left(\frac{1}{2}\right) = -\frac{27}{160} u^3\left(\frac{1}{2}\right) \\ u(0) = \frac{1}{2}u(1) + \frac{1}{100}u(\eta) + \frac{1}{100} \int_0^1 (u(s) - s)ds + \frac{1}{150} & \eta \in [0, 1]. \end{cases} \quad (4.1)$$

Let  $f(t, x, y, z, w) = \frac{t^4 x}{100} - \frac{t}{600}y - \frac{1}{100}z - t^3 w$ ,  $M(t) = 0$ ,  $N(t) = \frac{t}{600}$ ,  $K(t) = \frac{1}{100}$ ,  $H(t) = t^3$ ,  $k(t, s) = ts$ ,  $h(t, s) = \frac{1}{1000}$ ,  $Tu(t) = t \int_0^t su(s)ds$ ,  $Su(t) = \int_0^1 \frac{1}{1000}u(s)ds$ ,  $\alpha(t) = \frac{t}{2}$ ,  $\beta(t) = t$ ,  $\gamma(s) = s$ ,  $\delta(s) = s$ ,  $w(s, u(s)) = u(s) - s$ .

We can easily verify that (4.1) admits the lower solution  $u_0(t) = 0$  and the upper solution

$$v_0(t) = \begin{cases} \frac{2}{3}t + 1, & t \in \left[0, \frac{1}{2}\right], \\ \frac{2}{3}t + \frac{2}{3}, & t \in \left(\frac{1}{2}, 1\right], \end{cases}$$

and that  $u_0(t) \leq v_0(t)$ . It is easy to see that

$$\begin{aligned} I_k(x(t_k)) - I_k(y(t_k)) &= -\frac{27}{160}(x^3(t_k) - y^3(t_k)) \\ &\geq -\frac{3}{10}(x(t_k) - y(t_k)) \\ &= -L_1(x(t_k) - y(t_k)), \end{aligned}$$

where  $u_0(t_k) \leq y(t_k) \leq x(t_k) \leq v_0(t_k)$ ,  $L_1 = \frac{3}{10}$ .

Obviously,

$$\begin{aligned} f(t, \bar{u}, \bar{u}(\alpha(t)), T\bar{u}, S\bar{u}) - f(t, u, u(\alpha(t)), Tu, Su) &\geq -M(t)(\bar{u} - u) - N(t)(\bar{u} - u)(\alpha(t)) \\ &\quad - K(t)T(\bar{u} - u) - H(t)S(\bar{u} - u), \end{aligned}$$

$$W(t, \bar{u}(t)) - W(t, u(t)) = \bar{u}(t) - u(t) \geq \frac{t}{3}(\bar{u}(t) - u(t)),$$

for all  $u_0(t) \leq u(t) \leq \bar{u}(t) \leq v_0(t)$  in  $J$ .

And we can check that  $r^* = r = \frac{1}{2}$ ,  $[\mu^* + \sum_{k=1}^m L_k] < r^*$ ,  $e^{\int_0^T |M(\tau)|d\tau} (1 + \frac{r}{e^{\int_0^T M(\tau)d\tau} - r})(\mu + \sum_{k=1}^m L_k) < 0.91 < 1$ . Then all conditions of Theorem 3.1 are satisfied. Therefore, the conclusion of Theorem 3.1 holds for problem (4.1).

#### References

- [1] Meiqiang Feng, Dongxiu Xie, Multiple positive solutions of multi-point boundary value problem for second-order impulsive differential equations, J. Comput. Appl. Math. 223 (2009) 438–448.
- [2] Xiaoming He, Weigao Ge, Triple solutions for second-order three-point boundary value problems, J. Math. Anal. Appl. 268 (2002) 256–265.
- [3] Bing Liu, Jianshe Yu, Existence of solution for  $m$ -point boundary value problems of second-order differential systems with impulses, Appl. Math. Comput. 125 (2002) 155–175.
- [4] JuanJuan Xu, Ping Kang, Zhongli Wei, Singular multipoint impulsive boundary value problem with  $P$ -Laplacian operator, J. Appl. Math. Comput. 30 (2009) 105–120.
- [5] Tadeusz Jankowski, Monotone and numerical-analytic methods for differential equations, Comput. Math. Appl. 45 (2003) 1823–1828.
- [6] Guoping Chen, Jianhua Shen, Integral boundary value problems for first order impulsive functional differential equations, Int. J. Math. Anal. 1 (2007) 965–974.
- [7] Wei Din, Maoan Han, Junrong Mi, Periodic boundary value problems for the second order impulsive functional equations, Appl. Math. Comput. 50 (2005) 491–507.
- [8] Zhimin He, Jianshe Yu, Periodic boundary value problem for first order impulsive ordinary differential equation, J. Math. Anal. Appl. 272 (2002) 67–78.
- [9] Z. He, X. He, Monotone iterative technique for impulsive integro-differential equations with periodic boundary conditions, Comput. Math. Appl. 48 (2004) 73–84.
- [10] Jianli Li, Jianhua Shen, Periodic boundary value problems for impulsive integro-differential equations, Appl. Math. Comput. 183 (2006) 890–902.
- [11] Zhiguo Luo, Juan J. Nieto, New results for the periodic boundary value problem for impulsive integro-differential equations, Nonlinear Anal. 70 (2009) 2248–2260.
- [12] Zhenhai Liu, Anti-periodic solutions to nonlinear evolution equations, J. Funct. Anal. 258 (6) (2010) 2026–2033.
- [13] Jinbo Liu, Zhenhai Liu, On the existence of anti-periodic solutions for implicit differential equations, Acta Math. Hungar. 132 (3) (2011) 294–305.

- [14] Zhiguo Luo, Jianhua Shen, J.J. Nieto, Anti-periodic boundary value problem for first order impulsive ordinary differential equations, *Comput. Math. Appl.* 49 (2005) 253–261.
- [15] Xiaohuan Wang, Jihui Zhang, Impulsive anti-periodic boundary value problem of first-order integro-differential equations, *Comput. Math. Appl.* (2010) <http://dx.doi.org/10.1016/j.cam.2010.04.024>.
- [16] Tadeusz Jankowski, Boundary value problems for first order differential equations of mixed type, *Nonlinear Anal.* 64 (2006) 1984–1997.
- [17] Yong-Hoon Lee, Xinzhi Liu, Study of singular boundary value problems for second order impulsive differential equations, *Nonlinear Anal.* 70 (2009) 2736–2751.
- [18] J.J. Nieto, R. Rodríguez-López, New comparison results for impulsive integro-differential equations and applications, *J. Math. Anal. Appl.* 328 (2007) 1343–1368.
- [19] Yian Zhao, Guangxing Song, Xiaoyan Sun, Integral boundary value problems with causal operators, *Comput. Math. Appl.* 59 (2010) 2768–2775.
- [20] Dajun Guo, V. Lakshmikantham, Xinzhi Liu, *Nonlinear Integral Equations in Abstract Spaces*, Kluwer Academic Publishers, 1996.
- [21] V. Lakshmikantham, D.D. Bainov, P.S. Simeonov, *Theory of Impulsive Differential Equations*, World Scientific, Singapore, 1989.